arXiv:1702.08667v1 [math.NT] 28 Feb 2017

THETA SUMS OF HIGHER INDEX

JAE-HYUN YANG

Abstract. In this paper, we obtain some behaviours of theta sums of higher index for the Schrödinger-Weil representation of the Jacobi group associated with a positive definite symmetric real matrix of degree m.

1. Introduction

For a given fixed positive integer n, we let

$$\mathbb{H}_n = \left\{ \Omega \in \mathbb{C}^{(n,n)} \,\middle|\, \Omega = {}^t\Omega, \quad \operatorname{Im}\Omega > 0 \right\}$$

be the Siegel upper half plane of degree n and let

$$Sp(n,\mathbb{R}) = \left\{ g \in \mathbb{R}^{(2n,2n)} \,\middle|\, {}^t g J_n g = J_n \right\}$$

be the symplectic group of degree n, where $F^{(k,l)}$ denotes the set of all $k \times l$ matrices with entries in a commutative ring F for two positive integers k and l, ${}^tM$ denotes the transpose of a matrix M, $\operatorname{Im}\Omega$ denotes the imaginary part of Ω and

$$J_n = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}.$$

Here I_n denotes the identity matrix of degree n. We see that $Sp(n,\mathbb{R})$ acts on $\mathbb{H}_n$ transitively by

$$g \cdot \Omega = (A\Omega + B)(C\Omega + D)^{-1},$$

where $g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(n,\mathbb{R})$ and $\Omega \in \mathbb{H}_n$.

For two positive integers n and m, we consider the Heisenberg group

$$H_{\mathbb{R}}^{(n,m)} = \{\, (\lambda,\mu;\kappa) \mid \lambda,\mu \in \mathbb{R}^{(m,n)},\ \kappa \in \mathbb{R}^{(m,m)},\ \ \kappa + \mu\,{}^t\lambda \text{ symmetric} \,\}$$

endowed with the following multiplication law

$$(\lambda,\mu;\kappa) \circ (\lambda',\mu';\kappa') = (\lambda+\lambda', \mu+\mu'; \kappa+\kappa'+\lambda\,{}^t\mu' - \mu\,{}^t\lambda').$$

We let

$$G^J = Sp(n,\mathbb{R}) \ltimes H_{\mathbb{R}}^{(n,m)} \qquad \text{(semi-direct product)}$$

be the Jacobi group endowed with the following multiplication law

$$\Big(g,(\lambda,\mu;\kappa)\Big) \cdot \Big(g',(\lambda',\mu';\kappa')\Big) = \Big(gg', (\widetilde{\lambda}+\lambda', \widetilde{\mu}+\mu'; \kappa+\kappa'+\widetilde{\lambda}\,{}^t\mu' - \widetilde{\mu}\,{}^t\lambda')\Big)$$

2010 Mathematics Subject Classification: Primary 11F27, 11F50

Keywords and phrases: the Schrödinger representation, the Schrödinger-Weil representation, theta sums.

The author was supported by Basic Science Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (49562-1) and also by INHA UNIVERSITY Research Grant.

with $g, g' \in Sp(n,\mathbb{R}), (\lambda,\mu;\kappa),\ (\lambda',\mu';\kappa') \in H_{\mathbb{R}}^{(n,m)}$ and $(\widetilde{\lambda},\widetilde{\mu}) = (\lambda,\mu)g'$. Then we have the *natural transitive action* of G^J on the Siegel-Jacobi space $\mathbb{H}_{n,m} := \mathbb{H}_n \times \mathbb{C}^{(m,n)}$ defined by

$$\Big(g,(\lambda,\mu;\kappa)\Big)\cdot(\Omega,Z) = \Big((A\Omega+B)(C\Omega+D)^{-1}, (Z+\lambda\,\Omega+\mu)(C\,\Omega+D)^{-1}\Big),$$

where $g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(n,\mathbb{R}),\ (\lambda,\mu;\kappa) \in H_{\mathbb{R}}^{(n,m)}$ and $(\Omega,Z) \in \mathbb{H}_{n,m}$. Thus $\mathbb{H}_{n,m}$ is a homogeneous Kähler space which is not symmetric. In fact, $\mathbb{H}_{n,m}$ is biholomorphic to the homogeneous space G^J/K^J, where $K^J \cong U(n)\times S(m,\mathbb{R})$. Here $U(n)$ denotes the unitary group of degree n and $S(m,\mathbb{R})$ denote the abelian additive group consisting of all $m\times m$ symmetric real matrices. We refer to [1, 2, 4], [17]-[29] for more details on materials related to the Siegel-Jacobi space, e.g., Jacobi forms, invariant metrics, invariant differential operators and Maass-Jacobi forms.

The Weil representation for a symplectic group was first introduced by A. Weil in [10] to reformulate Siegel's analytic theory of quadratic forms (cf. [9]) in terms of the group theoretical theory. It is well known that the Weil representation plays a central role in the study of the transformation behaviors of theta series. In [28], Yang constructed the Schrödinger-Weil representation $\omega_{\mathcal{M}}$ of the Jacobi group G^J associated with a positive definite symmetric real matrix $\mathcal{M}$ of degree n explicitly.

This paper is organized as follows. In Section 2, we review the Schrödinger-Weil representation $\omega_{\mathcal{M}}$ of the Jacobi group G^J associated with a symmetric positive definite matrix $\mathcal{M}$ and recall the basic actions of $\omega_{\mathcal{M}}$ on the representation space $L^2\big(\mathbb{R}^{(m,n)}\big)$ which were expressed explicitly in [28]. In Section 3, we define the theta sum $\Theta_f^{[\mathcal{M}]}(\tau,\phi\,;\lambda,\mu,\kappa)$ of higher index and obtain some properties of the theta sum. The theta sum $\Theta_f^{[\mathcal{M}]}(\tau,\phi\,;\lambda,\mu,\kappa)$ is a generalization of the theta sum defined by J. Marklof [6].

Notations: We denote by $\mathbb{Z}$, $\mathbb{R}$ and $\mathbb{C}$ the ring of integers, the field of real numbers and the field of complex numbers respectively. $\mathbb{C}^\times$ denotes the multiplicative group of nonzero complex numbers and $\mathbb{Z}^\times$ denotes the set of all nonzero integers. T denotes the multiplicative group of complex numbers of modulus one. The symbol ":=" means that the expression on the right is the definition of that on the left. For two positive integers k and l, $F^{(k,l)}$ denotes the set of all $k\times l$ matrices with entries in a commutative ring F. For a square matrix $A \in F^{(k,k)}$ of degree k, $\sigma(A)$ denotes the trace of A. For any $M \in F^{(k,l)}$, ${}^tM$ denotes the transpose of a matrix M. I_n denotes the identity matrix of degree n. We put $i = \sqrt{-1}$. For a positive integer m we denote by $S(m,F)$ the additive group consisting of all $m\times m$ symmetric matrices with coefficients in a commutative ring F.

2. The Schrödinger-Weil Representation

In this section we review the Schrödinger-Weil representation of the Jacobi group G^J (cf. [28], Section 3).

Throughout this section we assume that $\mathcal{M}$ is a positive definite symmetric real $m\times m$ matrix. We let

$$L = \Big\{\,(0,\mu;\kappa) \in H_{\mathbb{R}}^{(n,m)} \,\Big|\, \mu\in\mathbb{R}^{(m,n)},\ \kappa = {}^t\kappa \in \mathbb{R}^{(m,m)}\,\Big\}.$$

be a commutative normal subgroup of $H_{\mathbb{R}}^{(n,m)}$ and $\chi_{\mathcal{M}} : L \longrightarrow \mathbb{C}^x$ be the unitary character of L defined by

$$\chi_{\mathcal{M}}((0,\mu;\kappa)) := e^{\pi\, i\, \sigma(\mathcal{M}\kappa)}, \quad (0,\mu;\kappa) \in L.$$

The representation $\mathscr{W}_{\mathcal{M}} = \mathrm{Ind}_{L}^{H_{\mathbb{R}}^{(n,m)}} \chi_{\mathcal{M}}$ induced by $\chi_{\mathcal{M}}$ from L is realized on the Hilbert space $H(\chi_{\mathcal{M}}) \cong L^2\left(\mathbb{R}^{(m,n)}, d\xi\right)$. $\mathscr{W}_{\mathcal{M}}$ is irreducible (cf. [11], Theorem 3) and is called the Schrödinger representation $\mathscr{W}_{\mathcal{M}}$ of the Heisenberg group $H_{\mathbb{R}}^{(n,m)}$ with the central character $\chi_{\mathcal{M}}$. We refer to [11, 12, 13, 14, 15, 16] for more details on representations of the Heisenberg group $H_{\mathbb{R}}^{(n,m)}$ and their related topics. Then $\mathscr{W}_{\mathcal{M}}$ is expressed explicitly as

$$[\mathscr{W}_{\mathcal{M}}(h_0)f]\,(\lambda) = e^{\pi\, i\, \sigma\{\mathcal{M}(\kappa_0+\mu_0\,{}^t\lambda_0+2\lambda\,{}^t\mu_0)\}}\, f(\lambda+\lambda_0), \tag{2.1}$$

where $h_0 = (\lambda_0, \mu_0; \kappa_0) \in H$ and $\lambda \in \mathbb{R}^{(m,n)}$. See Formula (2.4) in [28] for more detail on $\mathscr{W}_{\mathcal{M}}$. We note that the symplectic group $Sp(n,\mathbb{R})$ acts on $H_{\mathbb{R}}^{(n,m)}$ by conjugation inside G^J. For a fixed element $g \in Sp(n,\mathbb{R})$, the irreducible unitary representation $\mathscr{W}_{\mathcal{M}}^{g}$ of $H_{\mathbb{R}}^{(n,m)}$ defined by

$$\mathscr{W}_{\mathcal{M}}^{g}(h) = \mathscr{W}_{\mathcal{M}}(ghg^{-1}), \quad h \in H_{\mathbb{R}}^{(n,m)} \tag{2.2}$$

has the property that

$$\mathscr{W}_{\mathcal{M}}^{g}((0,0;\kappa)) = \mathscr{W}_{\mathcal{M}}((0,0;\kappa)) = e^{\pi i\, \sigma(\mathcal{M}\kappa)}\, \mathrm{Id}_{H(\chi_{\mathcal{M}})}, \quad \kappa \in S(m,\mathbb{R}).$$

Here $\mathrm{Id}_{H(\chi_{\mathcal{M}})}$ denotes the identity operator on the Hilbert space $H(\chi_{\mathcal{M}})$. According to Stone-von Neumann theorem, there exists a unitary operator $R_{\mathcal{M}}(g)$ on $H(\chi_{\mathcal{M}})$ with $R_{\mathcal{M}}(I_{2n}) = \mathrm{Id}_{H(\chi_{\mathcal{M}})}$ such that

$$R_{\mathcal{M}}(g)\mathscr{W}_{\mathcal{M}}(h) = \mathscr{W}_{\mathcal{M}}^{g}(h) R_{\mathcal{M}}(g) \qquad \text{for all } h \in H_{\mathbb{R}}^{(n,m)}. \tag{2.3}$$

We observe that $R_{\mathcal{M}}(g)$ is determined uniquely up to a scalar of modulus one.

From now on, for brevity, we put $G = Sp(n,\mathbb{R})$. According to Schur's lemma, we have a map $c_{\mathcal{M}} : G \times G \longrightarrow T$ satisfying the relation

$$R_{\mathcal{M}}(g_1 g_2) = c_{\mathcal{M}}(g_1, g_2) R_{\mathcal{M}}(g_1) R_{\mathcal{M}}(g_2) \quad \text{for all } \ g_1, g_2 \in G. \tag{2.4}$$

We recall that T denotes the multiplicative group of complex numbers of modulus one. Therefore $R_{\mathcal{M}}$ is a projective representation of G on $H(\chi_{\mathcal{M}})$ and $c_{\mathcal{M}}$ defines the cocycle class in $H^2(G,T)$. The cocycle $c_{\mathcal{M}}$ yields the central extension $G_{\mathcal{M}}$ of G by T. The group $G_{\mathcal{M}}$ is a set $G \times T$ equipped with the following multiplication

$$(g_1,t_1)\cdot(g_2,t_2) = \left(g_1 g_2, t_1 t_2\, c_{\mathcal{M}}(g_1,g_2)^{-1}\right), \quad g_1, g_2 \in G,\ t_1, t_2 \in T. \tag{2.5}$$

We see immediately that the map $\widetilde{R}_{\mathcal{M}} : G_{\mathcal{M}} \longrightarrow GL(H(\chi_{\mathcal{M}}))$ defined by

$$\widetilde{R}_{\mathcal{M}}(g,t) = t\, R_{\mathcal{M}}(g) \quad \text{for all } (g,t) \in G_{\mathcal{M}} \tag{2.6}$$

is a *true* representation of $G_{\mathcal{M}}$. As in Section 1.7 in [5], we can define the map $s_{\mathcal{M}} : G \longrightarrow T$ satisfying the relation

$$c_{\mathcal{M}}(g_1,g_2)^2 = s_{\mathcal{M}}(g_1)^{-1} s_{\mathcal{M}}(g_2)^{-1} s_{\mathcal{M}}(g_1 g_2) \quad \text{for all } g_1, g_2 \in G.$$

Thus we see that

$$G_{2,\mathcal{M}} = \left\{ (g,t) \in G_{\mathcal{M}} \mid t^2 = s_{\mathcal{M}}(g)^{-1} \right\} \tag{2.7}$$

is the metaplectic group associated with $\mathcal{M}$ that is a two-fold covering group of G. The restriction $R_{2,\mathcal{M}}$ of $\widetilde{R}_{\mathcal{M}}$ to $G_{2,\mathcal{M}}$ is the Weil representation of G associated with $\mathcal{M}$.

If we identify $h = (\lambda, \mu; \kappa) \in H_{\mathbb{R}}^{(n,m)}$ (resp. $g \in Sp(n,\mathbb{R})$) with $(I_{2n}, (\lambda, \mu; \kappa)) \in G^J$ (resp. $(g, (0,0;0)) \in G^J$), every element $\tilde{g}$ of G^J can be written as $\tilde{g} = hg$ with $h \in H_{\mathbb{R}}^{(n,m)}$ and $g \in Sp(n,\mathbb{R})$. In fact,

$$(g, (\lambda, \mu; \kappa)) = (I_{2n}, ((\lambda, \mu)g^{-1}; \kappa))\,(g, (0,0;0)) = ((\lambda, \mu)g^{-1}; \kappa) \cdot g.$$

Therefore we define the *projective* representation $\pi_{\mathcal{M}}$ of the Jacobi group G^J with cocycle $c_{\mathcal{M}}(g_1, g_2)$ by

$$\pi_{\mathcal{M}}(hg) = \mathscr{W}_{\mathcal{M}}(h)\, R_{\mathcal{M}}(g), \quad h \in H_{\mathbb{R}}^{(n,m)},\ g \in G. \tag{2.8}$$

We let

$$G_{\mathcal{M}}^J = G_{\mathcal{M}} \ltimes H_{\mathbb{R}}^{(n,m)}$$

be the semidirect product of $G_{\mathcal{M}}$ and $H_{\mathbb{R}}^{(n,m)}$ with the multiplication law

$$\begin{aligned} & \big((g_1,t_1),(\lambda_1,\mu_1;\kappa_1)\big) \cdot \big((g_2,t_2),(\lambda_2,\mu_2\,;\kappa_2)\big) \\ = \ & \big((g_1,t_1)(g_2,t_2), (\tilde{\lambda}+\lambda_2, \tilde{\mu}+\mu_2\,;\kappa_1+\kappa_2+\tilde{\lambda}\,{}^t\mu_2 - \tilde{\mu}\,{}^t\lambda_2)\big), \end{aligned}$$

where $(g_1,t_1),(g_2,t_2) \in G_{\mathcal{M}}$, $(\lambda_1,\mu_1;\kappa_1),(\lambda_2,\mu_2\,;\kappa_2) \in H_{\mathbb{R}}^{(n,m)}$ and $(\tilde{\lambda},\tilde{\mu}) = (\lambda,\mu)g_2$. If we identify $h = (\lambda,\mu\,;\kappa) \in H_{\mathbb{R}}^{(n,m)}$ (resp. $(g,t) \in G_{\mathcal{M}}$) with $((I_{2n},1),(\lambda,\mu\,;\kappa)) \in G_{\mathcal{M}}^J$ (resp. $((g,t),(0,0;0)) \in G_{\mathcal{M}}^J$), we see easily that every element $\big((g,t),(\lambda,\mu\,;\kappa)\big)$ of $G_{\mathcal{M}}^J$ can be expressed as

$$\big((g,t),(\lambda,\mu\,;\kappa)\big) = \big((I_{2n},1),((\lambda,\mu)g^{-1};\kappa)\big)\big((g,t),(0,0;0)\big) = ((\lambda,\mu)g^{-1};\kappa)(g,t).$$

Now we can define the *true* representation $\widetilde{\omega}_{\mathcal{M}}$ of $G_{\mathcal{M}}^J$ by

$$\widetilde{\omega}_{\mathcal{M}}(h\cdot(g,t)) = t\,\pi_{\mathcal{M}}(hg) = t\,\mathscr{W}_{\mathcal{M}}(h)\, R_{\mathcal{M}}(g), \quad h \in H_{\mathbb{R}}^{(n,m)},\ (g,t) \in G_{\mathcal{M}}. \tag{2.9}$$

We recall that the following matrices

$$\begin{aligned} t(b) &= \begin{pmatrix} I_n & b \\ 0 & I_n \end{pmatrix} \text{ with any } b = {}^tb \in \mathbb{R}^{(n,n)}, \\ g(\alpha) &= \begin{pmatrix} {}^t\alpha & 0 \\ 0 & \alpha^{-1} \end{pmatrix} \text{ with any } \alpha \in GL(n,\mathbb{R}), \\ \sigma_n &= \begin{pmatrix} 0 & -I_n \\ I_n & 0 \end{pmatrix} \end{aligned}$$

generate the symplectic group $G = Sp(n,\mathbb{R})$ (cf. [3, p. 326], [7, p. 210]). Therefore the following elements $h_t(\lambda,\mu\,;\kappa)$, $t(b\,;t)$, $g(\alpha\,;t)$ and $\sigma_{n\,;t}$ of $G_{\mathcal{M}} \ltimes H_{\mathbb{R}}^{(n,m)}$ defined by

$$\begin{aligned} & h_t(\lambda,\mu\,;\kappa) = \big((I_{2n},t),(\lambda,\mu;\kappa)\big) \text{ with } t \in T,\ \lambda,\mu \in \mathbb{R}^{(m,n)} \text{ and } \kappa \in \mathbb{R}^{(m,m)}, \\ & t(b\,;t) = \big((t(b),t),(0,0;0)\big) \text{ with any } b = {}^tb \in \mathbb{R}^{(n,n)},\ t \in T, \\ & g(\alpha\,;t) = \big(\big(g(\alpha),t\big),(0,0;0)\big) \text{ with any } \alpha \in GL(n,\mathbb{R}) \text{ and } t \in T, \\ & \sigma_{n\,;t} = ((\sigma_n,t),(0,0;0)) \text{ with } t \in T \end{aligned}$$

generate the group $G_{\mathcal{M}} \ltimes H_{\mathbb{R}}^{(n,m)}$. We can show that the representation $\widetilde{\omega}_{\mathcal{M}}$ is realized on the representation $H(\chi_{\mathcal{M}}) = L^2\big(\mathbb{R}^{(m,n)}\big)$ as follows: for each $f \in L^2\big(\mathbb{R}^{(m,n)}\big)$ and $x \in \mathbb{R}^{(m,n)}$, the actions of $\widetilde{\omega}_{\mathcal{M}}$ on the generators are given by

$$\big[\widetilde{\omega}_{\mathcal{M}}\big(h_t(\lambda,\mu\,;\kappa)\big)f\big]\,(x) \;=\; t\,e^{\pi i\,\sigma\{\mathcal{M}(\kappa+\mu\,{}^t\lambda+2\,x\,{}^t\mu)\}}\,f(x+\lambda), \tag{2.10}$$

$$\big[\widetilde{\omega}_{\mathcal{M}}\big(t(b\,;t)\big)f\big]\,(x) \;=\; t\,e^{\pi i\,\sigma(\mathcal{M}\,x\,b\,{}^t x)}f(x), \tag{2.11}$$

$$\big[\widetilde{\omega}_{\mathcal{M}}\big(g(\alpha\,;t)\big)f\big]\,(x) \;=\; t\,|\det\alpha|^{\frac{m}{2}}\,f(x\,{}^t\alpha), \tag{2.12}$$

$$\big[\widetilde{\omega}_{\mathcal{M}}\big(\sigma_{n\,;t}\big)f\big]\,(x) = t\,\big(\det\mathcal{M}\big)^{\frac{n}{2}}\int_{\mathbb{R}^{(m,n)}} f(y)\,e^{-2\,\pi i\,\sigma(\mathcal{M}\,y\,{}^t x)}\,dy. \tag{2.13}$$

Let

$$G_{2,\mathcal{M}}^{J} = G_{2,\mathcal{M}} \ltimes H_{\mathbb{R}}^{(n,m)}$$

be the semidirect product of $G_{2,\mathcal{M}}$ and $H_{\mathbb{R}}^{(n,m)}$. Then $G_{2,\mathcal{M}}^{J}$ is a subgroup of $G_{\mathcal{M}}^{J}$ which is a two-fold covering group of the Jacobi group G^J. The restriction $\omega_{\mathcal{M}}$ of $\widetilde{\omega}_{\mathcal{M}}$ to $G_{2,\mathcal{M}}^{J}$ is called the Schrödinger-Weil representation of G^J associated with $\mathcal{M}$.

Remark 2.1. *In the case $n = m = 1$, $\omega_{\mathcal{M}}$ is dealt in* [1] *and* [6].

Remark 2.2. *The Schrödinger-Weil representation is applied usefully to the theory of Maass-Jacobi forms* [8].

3. Theta Sums of Higher Index

Let $\mathcal{M}$ be a positive definite symmetric real matrix of degree m. We recall the Schrödinger representation $\mathscr{W}_{\mathcal{M}}$ of the Heisenberg group $H_{\mathbb{R}}^{(n,m)}$ associated with $\mathcal{M}$ that is given by Formula (2.1) in Section 2. We note that for an element $(\lambda,\mu;\kappa)$ of $H_{\mathbb{R}}^{(n,m)}$, we have the decomposition

$$(\lambda,\mu;\kappa) = (\lambda,0;0)\circ(0,\mu;0)\circ(0,0;\kappa-\lambda\,{}^t\mu).$$

We consider the embedding $\Phi_n : SL(2,\mathbb{R}) \longrightarrow Sp(n,\mathbb{R})$ defined by

$$\Phi_n\left(\begin{pmatrix} a & b\\ c & d\end{pmatrix}\right) := \begin{pmatrix} aI_n & bI_n\\ cI_n & dI_n\end{pmatrix}, \qquad \begin{pmatrix} a & b\\ c & d\end{pmatrix} \in SL(2,\mathbb{R}). \tag{3.1}$$

For $x, y \in \mathbb{R}^{(m,n)}$, we put

$$(x,y)_{\mathcal{M}} := \sigma({}^t x\mathcal{M}y) \qquad \text{and} \qquad \|x\|_{\mathcal{M}} := \sqrt{(x,x)_{\mathcal{M}}}.$$

According to Formulas (2.11)-(2.13), for any $M = \begin{pmatrix} a & b\\ c & d\end{pmatrix} \in SL(2,\mathbb{R}) \hookrightarrow Sp(n,\mathbb{R})$ and $f \in L^2\left(\mathbb{R}^{(m,n)}\right)$, we have the following explicit representation

$$[R_{\mathcal{M}}(M)f](x) = \begin{cases} |a|^{\frac{mn}{2}} e^{ab\|x\|^2_{\mathcal{M}}\pi i} f(ax) & \text{if } c = 0,\\ (\det\mathcal{M})^{\frac{n}{2}}\,|c|^{-\frac{mn}{2}}\int_{\mathbb{R}^{(m,n)}} e^{\frac{\alpha(M,x,y,\mathcal{M})}{c}\pi i} f(y)dy & \text{if } c \neq 0,\end{cases} \tag{3.2}$$

where

$$\alpha(M,x,y,\mathcal{M}) = a\,\|x\|^2_{\mathcal{M}} + d\,\|y\|^2_{\mathcal{M}} - 2(x,y)_{\mathcal{M}}.$$

Indeed, if $a=0$ and $c\neq 0$, using the decomposition

$$M = \begin{pmatrix} 0 & -c^{-1} \\ c & d \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} c & d \\ 0 & c^{-1} \end{pmatrix}$$

and if $a\neq 0$ and $c\neq 0$, using the decomposition

$$M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a & c^{-1} \\ 0 & a^{-1} \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} ac & ad \\ 0 & (ac)^{-1} \end{pmatrix},$$

we obtain Formula (3.2).

If

$$M_1 = \begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix}, \quad M_2 = \begin{pmatrix} a_2 & b_2 \\ c_2 & d_2 \end{pmatrix} \quad \text{and} \quad M_3 = \begin{pmatrix} a_3 & b_3 \\ c_3 & d_3 \end{pmatrix} \in SL(2,\mathbb{R})$$

with $M_3 = M_1 M_2$, the corresponding cocycle is given by

$$c_{\mathcal{M}}(M_1,M_2) = e^{-i\,\pi\, mn\, \mathrm{sign}(c_1c_2c_3)/4}, \tag{3.3}$$

where

$$\mathrm{sign}(x) = \begin{cases} -1 & (x<0) \\ 0 & (x=0) \\ 1 & (x>0). \end{cases}$$

In the special case when

$$M_1 = \begin{pmatrix} \cos\phi_1 & -\sin\phi_1 \\ \sin\phi_1 & \cos\phi_1 \end{pmatrix} \quad \text{and} \quad M_2 = \begin{pmatrix} \cos\phi_2 & -\sin\phi_2 \\ \sin\phi_2 & \cos\phi_2 \end{pmatrix},$$

we find

$$c_{\mathcal{M}}(M_1,M_2) = e^{-i\,\pi\, mn\,(\sigma_{\phi_1}+\sigma_{\phi_2}-\sigma_{\phi_1+\phi_2})/4},$$

where

$$\sigma_\phi = \begin{cases} 2\nu & \text{if } \phi = \nu\pi \\ 2\nu+1 & \text{if } \nu\pi < \phi < (\nu+1)\pi. \end{cases}$$

It is well known that every $M\in SL(2,\mathbb{R})$ admits the unique Iwasawa decomposition

$$M = \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix} \begin{pmatrix} v^{1/2} & 0 \\ 0 & v^{-1/2} \end{pmatrix} \begin{pmatrix} \cos\phi & -\sin\phi \\ \sin\phi & \cos\phi \end{pmatrix}, \tag{3.4}$$

where $\tau = u+iv \in \mathbb{H}_1$ and $\phi\in[0,2\pi)$. This parametrization $M=(\tau,\phi)$ in $SL(2,\mathbb{R})$ leads to the natural action of $SL(2,\mathbb{R})$ on $\mathbb{H}_1\times[0,2\pi)$ defined by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}(\tau,\phi) := \left(\frac{a\tau+b}{c\tau+d},\ \phi + \arg(c\tau+d) \bmod 2\pi \right). \tag{3.5}$$

Lemma 3.1. *For two elements g_1 and g_2 in $SL(2,\mathbb{R})$, we let*

$$g_1 = \begin{pmatrix} 1 & u_1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} v_1^{1/2} & 0 \\ 0 & v_1^{-1/2} \end{pmatrix} \begin{pmatrix} \cos\phi_1 & -\sin\phi_1 \\ \sin\phi_1 & \cos\phi_1 \end{pmatrix}$$

and

$$g_2 = \begin{pmatrix} 1 & u_2 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} v_2^{1/2} & 0 \\ 0 & v_2^{-1/2} \end{pmatrix} \begin{pmatrix} \cos\phi_2 & -\sin\phi_2 \\ \sin\phi_2 & \cos\phi_2 \end{pmatrix}$$

be the Iwasawa decompositions of g_1 and g_2 respectively, where $u_1, u_2 \in \mathbb{R}$, $v_1 > 0$, $v_2 > 0$ and $0 \leq \phi_1, \phi_2 < 2\pi$. Let

$$g_3 = g_1 g_2 = \begin{pmatrix} 1 & u_3 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} v_3^{1/2} & 0 \\ 0 & v_3^{-1/2} \end{pmatrix} \begin{pmatrix} \cos\phi_3 & -\sin\phi_3 \\ \sin\phi_3 & \cos\phi_3 \end{pmatrix}$$

be the Iwasawa decomposition of $g_3 = g_1 g_2$. Then we have

$$\begin{aligned} u_3 &= \frac{A}{(u_2 \sin\phi_1 + \cos\phi_1)^2 + (v_2 \sin\phi_1)^2}, \\ v_3 &= \frac{v_1 v_2}{(u_2 \sin\phi_1 + \cos\phi_1)^2 + (v_2 \sin\phi_1)^2} \end{aligned}$$

and

$$\phi_3 = tan^{-1}\left[\frac{(v_2 \cos\phi_2 + u_2 \sin\phi_2)\tan\phi_1 + \sin\phi_2}{(-v_2 \sin\phi_2 + u_2 \cos\phi_2)\tan\phi_1 + \cos\phi_2}\right],$$

where

$$\begin{aligned} A &= u_1(u_2 \sin\phi_1 + \cos\phi_1)^2 + (u_1 v_2 - v_1 u_2)\sin^2\phi_1 \\ &\quad + v_1 u_2 \cos^2\phi_1 + v_1(u_2^2 + v_2^2 - 1)\sin\phi_1 \cos\phi_1. \end{aligned}$$

Proof. If $g \in SL(2,\mathbb{R})$ has the unique Iwasawa decomposition (3.4), then we get the following

$$\begin{aligned} a &= v^{1/2}\cos\phi + uv^{-1/2}\sin\phi, \\ b &= -v^{1/2}\sin\phi + uv^{-1/2}\cos\phi, \\ c &= v^{-1/2}\sin\phi, \quad d = v^{-1/2}\cos\phi, \\ u &= (ac+bd)\left(c^2+d^2\right)^{-1}, \quad v = \left(c^2+d^2\right)^{-1}, \quad \tan\phi = \frac{c}{d}. \end{aligned}$$

We set

$$g_3 = g_1 g_2 = \begin{pmatrix} a_3 & b_3 \\ c_3 & d_3 \end{pmatrix}.$$

Since

$$u_3 = (a_3 c_3 + b_3 d_3)\left(c_3^2 + d_3^2\right)^{-1}, \quad v = \left(c_3^2 + d_3^2\right)^{-1}, \quad \tan\phi_3 = \frac{c_3}{d_3},$$

by an easy computation, we obtain the desired results. □

Now we use the new coordinates $(\tau = u + iv, \phi)$ with $\tau \in \mathbb{H}_1$ and $\phi \in [0, 2\pi)$ in $SL(2,\mathbb{R})$. According to Formulas (2.11)-(2.13), the projective representation $R_{\mathcal{M}}$ of $SL(2,\mathbb{R}) \hookrightarrow Sp(n,\mathbb{R})$ reads in these coordinates $(\tau = u + iv, \phi)$ as follows:

$$[R_{\mathcal{M}}(\tau,\phi)f](x) = v^{\frac{mn}{4}}\, e^{u\|x\|^2_{\mathcal{M}}\pi\, i}\, [R_{\mathcal{M}}(i,\phi)f]\left(v^{1/2}x\right), \tag{3.6}$$

where $f \in L^2\left(\mathbb{R}^{(m,n)}\right)$, $x \in \mathbb{R}^{(m,n)}$ and

$$\begin{aligned} &[R_{\mathcal{M}}(i,\phi)f](x) \\ &= \begin{cases} f(x) & \text{if } \phi \equiv 0 \bmod 2\pi, \\ f(-x) & \text{if } \phi \equiv \pi \bmod 2\pi, \\ (\det\mathcal{M})^{\frac{n}{2}}\, |\sin\phi|^{-\frac{mn}{2}} \int_{\mathbb{R}^{(m,n)}} e^{B(x,y,\phi,\mathcal{M})\pi i}\, f(y)dy & \text{if } \phi \not\equiv 0 \bmod \pi. \end{cases} \end{aligned} \tag{3.7}$$

Here

$$B(x,y,\phi,\mathcal{M}) = \frac{\left(\|x\|^2_{\mathcal{M}} + \|y\|^2_{\mathcal{M}}\right)\cos\phi - 2(x,y)_{\mathcal{M}}}{\sin\phi}.$$

Now we set

$$S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

We note that

$$\left[R_{\mathcal{M}}\left(i,\frac{\pi}{2}\right)f\right](x) = \left[R_{\mathcal{M}}(S)f\right](x) = (\det\mathcal{M})^{\frac{n}{2}}\int_{\mathbb{R}^{(m,n)}} f(y)\, e^{-2\,(x,y)_{\mathcal{M}}\,\pi\,i}\,dy \tag{3.8}$$

for $f \in L^2\left(\mathbb{R}^{(m,n)}\right)$.

Remark 3.1. *For Schwartz functions $f \in \mathscr{S}\left(\mathbb{R}^{(m,n)}\right)$, we have*

$$\lim_{\phi\longrightarrow 0\pm} |\sin\phi|^{-\frac{mn}{2}}\int_{\mathbb{R}^{(m,n)}} e^{B(x,y,\phi,\mathcal{M})\,\pi\,i}\,f(y)dy = e^{\pm i\,\pi\,mn/4}f(x) \neq f(x).$$

Therefore the projective representation $R_{\mathcal{M}}$ is not continuous at $\phi = \nu\pi\,(\nu\in\mathbb{Z})$ in general. If we set

$$\tilde{R}_{\mathcal{M}}(\tau,\phi) = e^{-i\,\pi\,mn\sigma_\phi/4}R_{\mathcal{M}}(\tau,\phi),$$

$\tilde{R}_{\mathcal{M}}$ corresponds to a unitary representation of the double cover of $SL(2,\mathbb{R})$ (cf. Formula (2.6) and [5]). This means in particular that

$$\tilde{R}_{\mathcal{M}}(i,\phi)\tilde{R}_{\mathcal{M}}(i,\phi') = \tilde{R}_{\mathcal{M}}(i,\phi+\phi'),$$

where $\phi\in[0,4\pi)$ parametrises the double cover of $SO(2)\subset SL(2,\mathbb{R})$.

We observe that for any element $(g,(\lambda,\mu;\kappa)) \in G^J$ with $g\in Sp(n,\mathbb{R})$ and $(\lambda,\mu;\kappa)\in H^{(n,m)}_{\mathbb{R}}$, we have the following decomposition

$$(g,(\lambda,\mu;\kappa)) = (I_{2n},((\lambda,\mu)g^{-1};\kappa))\,(g,(0,0;0)) = ((\lambda,\mu)g^{-1};\kappa)\cdot g.$$

Thus $Sp(n,\mathbb{R})$ acts on $H^{(n,m)}_{\mathbb{R}}$ naturally by

$$g\cdot(\lambda,\mu;\kappa) = \left((\lambda,\mu)g^{-1};\kappa\right), \qquad g\in Sp(n,\mathbb{R}),\ (\lambda,\mu;\kappa)\in H^{(n,m)}_{\mathbb{R}}.$$

Definition 3.1. *For any Schwartz function $f\in\mathscr{S}\left(\mathbb{R}^{(m,n)}\right)$, we define the function $\Theta^{[\mathcal{M}]}_f$ on the Jacobi group $SL(2,\mathbb{R})\ltimes H^{(n,m)}_{\mathbb{R}} \hookrightarrow G^J$ by*

$$\Theta^{[\mathcal{M}]}_f(\tau,\phi\,;\lambda,\mu,\kappa) := \sum_{\omega\in\mathbb{Z}^{(m,n)}}\left[\pi_{\mathcal{M}}\left((\lambda,\mu;\kappa)(\tau,\phi)\right)f\right](\omega), \tag{3.9}$$

where $(\tau,\phi)\in SL(2,\mathbb{R})$ and $(\lambda,\mu\,;\kappa)\in H^{(n,m)}_{\mathbb{R}}$. The function $\Theta^{[\mathcal{M}]}_f$ is called the theta sum of index $\mathcal{M}$ associated to a Schwartz function f. The projective representation $\pi_{\mathcal{M}}$ of the Jacobi group G^J was already defined by Formula (2.8). More precisely, for $\tau = u+iv\in\mathbb{H}_1$ and $(\lambda,\mu;\kappa)\in H^{(n,m)}_{\mathbb{R}}$, we have

$$\begin{aligned}\Theta^{[\mathcal{M}]}_f(\tau,\phi\,;\lambda,\mu,\kappa) &= v^{\frac{mn}{4}}\ e^{2\,\pi\,i\,\sigma(\mathcal{M}(\kappa+\mu^t\lambda))} \\ &\times\sum_{\omega\in\mathbb{Z}^{(m,n)}} e^{\pi\,i\left\{u\|\omega+\lambda\|^2_{\mathcal{M}}+2(\omega,\mu)_{\mathcal{M}}\right\}}\ \left[R_{\mathcal{M}}(i,\phi)f\right]\left(v^{1/2}(\omega+\lambda)\right).\end{aligned}$$

Lemma 3.2. *We set $f_\phi := \tilde{R}_{\mathcal{M}}(i,\phi)f$ for $f \in \mathscr{S}\left(\mathbb{R}^{(m,n)}\right)$. Then for any $R > 1$, there exists a constant C_R such that for all $x \in \mathbb{R}^{(m,n)}$ and $\phi \in \mathbb{R}$,*

$$|f_\phi(x)| \le C_R\,(1 + \|x\|_{\mathcal{M}})^{-R}.$$

Proof. Following the arguments in the proof of Lemma 4.3 in [6], pp. 428-429, we get the desired result. □

Theorem 3.1 (Jacobi 1)**.** *Let $\mathcal{M}$ be a positive definite symmetric integral matrix of degree m such that $\mathcal{M}\mathbb{Z}^{(m,n)} = \mathbb{Z}^{(m,n)}$. Then for any Schwartz function $f \in \mathscr{S}\left(\mathbb{R}^{(m,n)}\right)$, we have*

$$\Theta_f^{[\mathcal{M}]}\left(-\frac{1}{\tau},\, \phi + arg\,\tau\,;\, -\mu, \lambda, \kappa\right) = \left(\det \mathcal{M}\right)^{-\frac{n}{2}} c_{\mathcal{M}}(S,(\tau,\phi))\,\Theta_f^{[\mathcal{M}]}(\tau,\phi\,;\lambda,\mu,\kappa),$$

where

$$c_{\mathcal{M}}(S,(\tau,\phi)) := e^{i\,\pi mn\, sign(\sin\phi\,\sin(\phi+\arg\tau))}.$$

Proof. First we recall that for any Schwartz function $\varphi \in \mathscr{S}\left(\mathbb{R}^{(m,n)}\right)$, the Fourier transform $\mathscr{F}\varphi$ of φ is given by

$$\left(\mathscr{F}\varphi\right)(x) = \int_{\mathbb{R}^{(m,n)}} \varphi(y)\, e^{-2\pi i\,\sigma(y\,{}^t x)} dy.$$

Now we put

$$S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \in SL(2,\mathbb{Z}) \hookrightarrow Sp(n,\mathbb{R})$$

and for any $F \in \mathscr{S}\left(\mathbb{R}^{(m,n)}\right)$, we put

$$F_{\mathcal{M}}(x) := F(\mathcal{M}^{-1}x), \quad x \in \mathbb{R}^{(m,n)}.$$

According to Formula (2.13), for any $F \in \mathscr{S}\left(\mathbb{R}^{(m,n)}\right)$,

$$\begin{aligned}
\left[R_{\mathcal{M}}(S)F\right](x) &= \left(\det \mathcal{M}\right)^{\frac{n}{2}} \int_{\mathbb{R}^{(m,n)}} F(y)\, e^{-2\pi i\,\sigma(\mathcal{M}y\,{}^t x)} dy \\
&= \left(\det \mathcal{M}\right)^{-\frac{n}{2}} \int_{\mathbb{R}^{(m,n)}} F(\mathcal{M}^{-1}y)\, e^{-2\pi i\,\sigma(y\,{}^t x)} dy \\
&= \left(\det \mathcal{M}\right)^{-\frac{n}{2}} \int_{\mathbb{R}^{(m,n)}} F_{\mathcal{M}}(y)\, e^{-2\pi i\,\sigma(y\,{}^t x)} dy \\
&= \left(\det \mathcal{M}\right)^{-\frac{n}{2}} \left[\mathscr{F}F_{\mathcal{M}}\right](x).
\end{aligned}$$

Thus we have

$$\mathscr{F}F_{\mathcal{M}} = \left(\det \mathcal{M}\right)^{\frac{n}{2}} R_{\mathcal{M}}(S)F \qquad \text{for } F \in \mathscr{S}\left(\mathbb{R}^{(m,n)}\right). \tag{3.10}$$

By Lemma 3.1, we get easily

$$S \cdot (\tau,\phi) = \left(-\frac{1}{\tau}, \phi + \arg\tau\right). \tag{3.11}$$

If we take $F = \pi_{\mathcal{M}}((\lambda,\mu\,;\kappa)(\tau,\phi))f$ for $f \in \mathscr{S}\left(\mathbb{R}^{(m,n)}\right)$, a fixed element $(\lambda,\mu\,;\kappa) \in H_{\mathbb{R}}^{(n,m)}$ and an fixed element $(\tau,\phi) \in SL(2,\mathbb{R})$, then it is easily seen that $F \in \mathscr{S}\left(\mathbb{R}^{(m,n)}\right)$.

According to Formulas (3.11), if we take $F = \pi_{\mathcal{M}}((\lambda,\mu\,;\kappa)(\tau,\phi))f$ for $f \in \mathscr{S}\left(\mathbb{R}^{(m,n)}\right)$,

$$\begin{aligned}
\big[R_{\mathcal{M}}(S)F\big](x) &= \big[R_{\mathcal{M}}(S)\pi_{\mathcal{M}}\big((\lambda,\mu\,;\kappa)(\tau,\phi)\big)f\big]\,(x), \quad x \in \mathbb{R}^{(m,n)} \\
&= \left[R_{\mathcal{M}}(S)\mathscr{W}_{\mathcal{M}}(\lambda,\mu\,;\kappa)R_{\mathcal{M}}(\tau,\phi)f\right](x) \\
&= \left[\mathscr{W}_{\mathcal{M}}\big((\lambda,\mu)S^{-1};\kappa\big)R_{\mathcal{M}}(S)R_{\mathcal{M}}(\tau,\phi)f\right](x) \\
&= c_{\mathcal{M}}(S,(\tau,\phi))^{-1}\,\left[\mathscr{W}_{\mathcal{M}}(-\mu,\lambda\,;\kappa)R_{\mathcal{M}}\big(S\cdot(\tau,\phi)\big)f\right](x) \\
&= c_{\mathcal{M}}(S,(\tau,\phi))^{-1}\,\left[\mathscr{W}_{\mathcal{M}}(-\mu,\lambda\,;\kappa)R_{\mathcal{M}}\left(-\frac{1}{\tau},\phi+\arg\tau\right)f\right](x) \\
&= c_{\mathcal{M}}(S,(\tau,\phi))^{-1}\,\left[\pi_{\mathcal{M}}\left((-\mu,\lambda\,;\kappa)\left(-\frac{1}{\tau},\phi+\arg\tau\right)\right)f\right](x).
\end{aligned}$$

Thus we obtain

$$\big[R_{\mathcal{M}}(S)F\big](x) = c_{\mathcal{M}}(S,(\tau,\phi))^{-1}\,\left[\pi_{\mathcal{M}}\left((-\mu,\lambda\,;\kappa)\left(-\frac{1}{\tau},\phi+\arg\tau\right)\right)f\right](x). \tag{3.12}$$

According to Poisson summation formula, we have

$$\sum_{\omega\in\mathbb{Z}^{(m,n)}}[\mathscr{F}F_{\mathcal{M}}]\,(\omega) = \sum_{\omega\in\mathbb{Z}^{(m,n)}}F_{\mathcal{M}}(\omega). \tag{3.13}$$

It follows from (3.10) and (3.12) that

$$\begin{aligned}
\sum_{\omega\in\mathbb{Z}^{(m,n)}}[\mathscr{F}F_{\mathcal{M}}]\,(\omega) &= \big(\det\mathcal{M}\big)^{\frac{n}{2}}\sum_{\omega\in\mathbb{Z}^{(m,n)}}\big[R_{\mathcal{M}}(S)F\big](\omega) \\
&= \big(\det\mathcal{M}\big)^{\frac{n}{2}}\,c_{\mathcal{M}}(S,(\tau,\phi))^{-1} \\
&\quad\times\sum_{\omega\in\mathbb{Z}^{(m,n)}}\left[\pi_{\mathcal{M}}\left((-\mu,\lambda\,;\kappa)\left(-\frac{1}{\tau},\phi+\arg\tau\right)\right)f\right](x) \\
&= \big(\det\mathcal{M}\big)^{\frac{n}{2}}\,c_{\mathcal{M}}(S,(\tau,\phi))^{-1}\,\Theta_f^{[\mathcal{M}]}\left(-\frac{1}{\tau},\,\phi+\arg\tau\,;-\mu,\lambda,\kappa\right).
\end{aligned}$$

On the other hand,

$$\begin{aligned}
\sum_{\omega\in\mathbb{Z}^{(m,n)}}F_{\mathcal{M}}(\omega) &= \sum_{\omega\in\mathbb{Z}^{(m,n)}}F\big(\mathcal{M}^{-1}\omega\big) \\
&= \sum_{\omega\in\mathbb{Z}^{(m,n)}}[\pi_{\mathcal{M}}((\lambda,\mu\,;\kappa)(\tau,\phi))f]\,\big(\mathcal{M}^{-1}\omega\big) \\
&= \sum_{\omega\in\mathbb{Z}^{(m,n)}}[\pi_{\mathcal{M}}((\lambda,\mu\,;\kappa)(\tau,\phi))f]\,(\omega) \quad \left(\because\ \mathcal{M}^{-1}\mathbb{Z}^{(m,n)} = \mathbb{Z}^{(m,n)}\right) \\
&= \Theta_f^{[\mathcal{M}]}(\tau,\phi\,;\lambda,\mu,\kappa).
\end{aligned}$$

Hence from (3.13) we obtain the desired formula

$$\Theta_f^{[\mathcal{M}]}\left(-\frac{1}{\tau},\,\phi+\arg\tau\,;-\mu,\lambda,\kappa\right) = \big(\det\mathcal{M}\big)^{-\frac{n}{2}}\,c_{\mathcal{M}}(S,(\tau,\phi))\,\Theta_f^{[\mathcal{M}]}(\tau,\phi\,;\lambda,\mu,\kappa).$$

If

$$S = \begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix}, \quad (\tau,\phi) = \begin{pmatrix} a_2 & b_2 \\ c_2 & d_2 \end{pmatrix} \quad \text{and} \quad S\cdot(\tau,\phi) = \begin{pmatrix} a_3 & b_3 \\ c_3 & d_3 \end{pmatrix} \in SL(2,\mathbb{R}),$$

according to Lemma 3.1, we get easily

$$c_1c_2c_3 = \left(u^2+v^2\right)^{1/2}\sin\phi\,\sin(\phi+\arg\tau),$$

where

$$(\tau,\phi) = \begin{pmatrix}1 & u\\ 0 & 1\end{pmatrix}\begin{pmatrix}v^{1/2} & 0\\ 0 & v^{-1/2}\end{pmatrix}\begin{pmatrix}\cos\phi & -\sin\phi\\ \sin\phi & \cos\phi\end{pmatrix}$$

is the Iwasawa decomposition of $(\tau,\phi)\in SL(2,\mathbb{R})$. Thus we obtain

$$c_{\mathcal{M}}(S,(\tau,\phi)) = e^{i\,\pi mn\,\mathrm{sign}(c_1c_2c_3)} = e^{i\,\pi mn\,\mathrm{sign}(\sin\phi\,\sin(\phi+\arg\tau))}.$$

This completes the proof. □

Theorem 3.2 (Jacobi 2)**.** *Let $\mathcal{M}=(\mathcal{M}_{kl})$ be a positive definite symmetric integral $m\times m$ matrix and let $s=(s_{kj})\in\mathbb{Z}^{(m,n)}$ be integral. Then we have*

$$\Theta_f^{[\mathcal{M}]}(\tau+2,\phi\,;\lambda,s-2\,\lambda+\mu,\kappa-s\,{}^t\lambda) = \Theta_f^{[\mathcal{M}]}(\tau,\phi\,;\lambda,\mu,\kappa)$$

for all $(\tau,\phi)\in SL(2,\mathbb{R})$ and $(\lambda,\mu;\kappa)\in H_{\mathbb{R}}^{(n,m)}$.

Proof. For brevity, we put $T_*=\begin{pmatrix}1 & 2\\ 0 & 1\end{pmatrix}$. According to Lemma 3.1, for any $(\tau,\phi)\in SL(2,\mathbb{R})$, the multiplication of T_* and (τ,ϕ) is given by

$$T_*(\tau,\phi) = (\tau+2,\phi). \tag{3.14}$$

For $s\in\mathbb{R}^{(m,n)}$, $(\lambda,\mu\,;\kappa)\in H_{\mathbb{R}}^{(n,m)}$ and $(\tau,\phi)\in SL(2,\mathbb{R})$, according to (3.14),

$$\begin{aligned}
&\pi_{\mathcal{M}}((0,s;0)T_*)\,\pi_{\mathcal{M}}((\lambda,\mu\,;\kappa)(\tau,\phi))\\
=\;& \mathscr{W}_{\mathcal{M}}(0,s;0)R_{\mathcal{M}}(T_*)\mathscr{W}_{\mathcal{M}}(\lambda,\mu\,;\kappa)R_{\mathcal{M}}(\tau,\phi)\\
=\;& \mathscr{W}_{\mathcal{M}}(0,s;0)\mathscr{W}_{\mathcal{M}}\big((\lambda,\mu)T_*^{-1}\,;\kappa\big)R_{\mathcal{M}}(T_*)R_{\mathcal{M}}(\tau,\phi)\\
=\;& c_{\mathcal{M}}(T_*,(\tau,\phi))^{-1}\mathscr{W}_{\mathcal{M}}(\lambda,s-2\,\lambda+\mu\,;\kappa-s\,{}^t\lambda)R_{\mathcal{M}}\big(T_*(\tau,\phi)\big)\\
=\;& \mathscr{W}_{\mathcal{M}}(\lambda,s-2\,\lambda+\mu\,;\kappa-s\,{}^t\lambda)R_{\mathcal{M}}(\tau+2,\phi)\\
=\;& \pi_{\mathcal{M}}\big((\lambda,s-2\,\lambda+\mu\,;\kappa-s\,{}^t\lambda)(\tau+2,\phi)\big).
\end{aligned}$$

Here we used the fact that $c_{\mathcal{M}}(T_*,(\tau,\phi))=1$ because T_* is upper triangular.

On the other hand, according to the assumptions on $\mathcal{M}$ and s, for $f\in\mathscr{S}\left(\mathbb{R}^{(m,n)}\right)$ and $\omega\in\mathbb{Z}^{(m,n)}$, using Formulas (2.1), (2.11) or (3.6), we have

$$\begin{aligned}
&\left[\pi_{\mathcal{M}}\big((0,s;0)T_*\big)\,\pi_{\mathcal{M}}\big((\lambda,\mu;\kappa)(\tau,\phi)\big)f\right](\omega)\\
=\;& \left[\mathscr{W}_{\mathcal{M}}(0,s;0)R_{\mathcal{M}}(T_*)\,\pi_{\mathcal{M}}\big((\lambda,\mu;\kappa)(\tau,\phi)\big)f\right](\omega)\\
=\;& e^{2\pi i\,\sigma(\mathcal{M}\omega\,{}^ts)}\cdot e^{2\,\|\omega\|^2_{\mathcal{M}}\pi\,i}\,\left[R_{\mathcal{M}}(i,0)\,\pi_{\mathcal{M}}\big((\lambda,\mu;\kappa)(\tau,\phi)\big)f\right](\omega)\\
=\;& \left[\pi_{\mathcal{M}}\big((\lambda,\mu;\kappa)(\tau,\phi)\big)f\right](\omega).
\end{aligned}$$

Here we used the facts that

$$e^{2\pi i\,\sigma(\mathcal{M}\omega\,{}^ts)}=1,\quad e^{2\,\|\omega\|^2_{\mathcal{M}}\pi\,i}=1\quad\text{and}\quad R_{\mathcal{M}}(i,0)f=f\ \text{(cf. (3.7))}.$$

Therefore for $f \in \mathscr{S}\left(\mathbb{R}^{(m,n)}\right)$,

$$\begin{aligned}
& \Theta_f^{[\mathcal{M}]}(\tau+2,\phi\,;\lambda,s-2\,\lambda+\mu,\kappa-s\,{}^t\lambda) \\
= & \sum_{\omega\in\mathbb{Z}^{(m,n)}} \left[\pi_{\mathcal{M}}\big((\lambda,s-2\,\lambda+\mu,\kappa-s\,{}^t\lambda)(\tau+2,\phi)\big)f\right](\omega) \\
= & \sum_{\omega\in\mathbb{Z}^{(m,n)}} \left[\pi_{\mathcal{M}}\big((0,s;0)T_*\big)\,\pi_{\mathcal{M}}\big((\lambda,\mu;\kappa)(\tau,\phi)\big)f\right](\omega) \\
= & \sum_{\omega\in\mathbb{Z}^{(m,n)}} \left[\pi_{\mathcal{M}}\big((\lambda,\mu;\kappa)(\tau,\phi)\big)f\right](\omega) \\
= & \Theta_f^{[\mathcal{M}]}(\tau,\phi\,;\lambda,\mu,\kappa).
\end{aligned}$$

This completes the proof. □

Theorem 3.3 (Jacobi 3)**.** *Let $\mathcal{M}=(\mathcal{M}_{kl})$ be a positive definite symmetric integral $m\times m$ matrix and let $(\lambda_0,\mu_0;\kappa_0)\in H_{\mathbb{Z}}^{(m,n)}$ be an integral element of $H_{\mathbb{R}}^{(n,m)}$. Then we have*

$$\begin{aligned}
& \Theta_f^{[\mathcal{M}]}(\tau,\phi\,;\lambda+\lambda_0,\mu+\mu_0,\kappa+\kappa_0+\lambda_0\,{}^t\mu-\mu_0\,{}^t\lambda) \\
= & \; e^{\pi\,i\,\sigma(\mathcal{M}(\kappa_0+\mu_0\,{}^t\lambda_0))}\Theta_f^{[\mathcal{M}]}(\tau,\phi\,;\lambda,\mu,\kappa)
\end{aligned}$$

for all $(\tau,\phi)\in SL(2,\mathbb{R})$ and $(\lambda,\mu;\kappa)\in H_{\mathbb{R}}^{(n,m)}$.

Proof. For any $f\in\mathscr{S}\left(\mathbb{R}^{(m,n)}\right)$, we have

$$\begin{aligned}
& \sum_{\omega\in\mathbb{Z}^{(m,n)}} \left[\mathscr{W}_{\mathcal{M}}(\lambda_0,\mu_0;\kappa_0)\pi_{\mathcal{M}}\big((\lambda,\mu\,;\kappa)(\tau,\phi)\big)f\right](\omega) \\
= & \sum_{\omega\in\mathbb{Z}^{(m,n)}} \left[\mathscr{W}_{\mathcal{M}}(\lambda_0,\mu_0;\kappa_0)\mathscr{W}_{\mathcal{M}}(\lambda,\mu\,;\kappa)R_{\mathcal{M}}(\tau,\phi)f\right](\omega) \\
= & \sum_{\omega\in\mathbb{Z}^{(m,n)}} \left[\mathscr{W}_{\mathcal{M}}(\lambda_0+\lambda,\mu_0+\mu;\kappa_0+\kappa+\lambda_0\,{}^t\mu-\mu_0\,{}^t\lambda))R_{\mathcal{M}}(\tau,\phi)f\right](\omega) \\
= & \sum_{\omega\in\mathbb{Z}^{(m,n)}} \left[\pi_{\mathcal{M}}\big((\lambda_0+\lambda,\mu_0+\mu;\kappa_0+\kappa+\lambda_0\,{}^t\mu-\mu_0\,{}^t\lambda)(\tau,\phi)\big)f\right](\omega) \\
= & \; \Theta_f^{[\mathcal{M}]}(\tau,\phi\,;\lambda+\lambda_0,\mu+\mu_0,\kappa+\kappa_0+\lambda_0\,{}^t\mu-\mu_0\,{}^t\lambda).
\end{aligned}$$

On the other hand, for any $f \in \mathscr{S}\left(\mathbb{R}^{(m,n)}\right)$, we have

$$\begin{aligned}
&\sum_{\omega\in\mathbb{Z}^{(m,n)}} \left[\mathscr{W}_{\mathcal{M}}(\lambda_0,\mu_0;\kappa_0)\pi_{\mathcal{M}}\big((\lambda,\mu\,;\kappa)(\tau,\phi)\big)f\right](\omega)\\
=&\sum_{\omega\in\mathbb{Z}^{(m,n)}} e^{\pi i\sigma\{\mathcal{M}(\kappa_0+\mu_0\,{}^t\lambda_0+2\,\omega\,{}^t\mu_0)\}}\left[\pi_{\mathcal{M}}(\tau,\phi\,;\lambda,\mu,\kappa)f\right](\omega+\lambda_0)\\
=&\ e^{\pi i\sigma\{\mathcal{M}(\kappa_0+\mu_0\,{}^t\lambda_0\}}\sum_{\omega\in\mathbb{Z}^{(m,n)}}\left[\pi_{\mathcal{M}}(\tau,\phi\,;\lambda,\mu,\kappa)f\right](\omega+\lambda_0)\quad(\because\ \mu_0 \text{ is integral})\\
=&\ e^{\pi i\sigma\{\mathcal{M}(\kappa_0+\mu_0\,{}^t\lambda_0\}}\sum_{\omega\in\mathbb{Z}^{(m,n)}}\left[\pi_{\mathcal{M}}(\tau,\phi\,;\lambda,\mu,\kappa)f\right](\omega)\quad(\because\ \lambda_0 \text{ is integral})\\
=&\ e^{\pi i\sigma\{\mathcal{M}(\kappa_0+\mu_0\,{}^t\lambda_0\}}\,\Theta_f^{[\mathcal{M}]}(\tau,\phi\,;\lambda,\mu,\kappa).
\end{aligned}$$

Finally we obtain the desired result. □

We put $V(m,n)=\mathbb{R}^{(m,n)}\times\mathbb{R}^{(m,n)}$. Let

$$G^{(m,n)} := SL(2,\mathbb{R})\ltimes V(m,n)$$

be the group with the following multiplication law

$$(g_1,(\lambda_1,\mu_1))\cdot(g_2,(\lambda_2,\mu_2)) = (g_1g_2,(\lambda_1,\mu_1)g_2+(\lambda_2,\mu_2)), \tag{3.15}$$

where $g_1,g_2\in SL(2,\mathbb{R})$ and $\lambda_1,\lambda_2,\mu_1,\mu_2\in\mathbb{R}^{(m,n)}$.

We define

$$\Gamma^{(m,n)} := SL(2,\mathbb{Z})\ltimes H_{\mathbb{Z}}^{(n,m)}.$$

Then $\Gamma^{(m,n)}$ acts on $G^{(m,n)}$ naturally through the multiplication law (3.15).

Lemma 3.3. *$\Gamma^{(m,n)}$ is generated by the elements*

$$(S,(0,0)),\quad (T_\flat,(0,s))\quad \textit{and}\quad (I_2,(\lambda_0,\mu_0)),$$

where

$$S=\begin{pmatrix}0&-1\\1&0\end{pmatrix},\quad T_\flat=\begin{pmatrix}1&1\\0&1\end{pmatrix}\quad \textit{and } s,\lambda_0,\mu_0\in\mathbb{Z}^{(m,n)}.$$

Proof. Since $SL(2,\mathbb{Z})$ is generated by S and $T_\flat$, we get the desired result. □

We define

$$\begin{aligned}
&\Theta_f^{[\mathcal{M}]}(\tau,\phi;\lambda,\mu)\\
=&\ v^{\frac{mn}{4}}\sum_{\omega\in\mathbb{Z}^{(m,n)}} e^{\pi\,i\left\{u\|\omega+\lambda\|^2_{\mathcal{M}}+2(\omega,\mu)_{\mathcal{M}}\right\}}\left[R_{\mathcal{M}}(i,\phi)f\right]\left(v^{1/2}(\omega+\lambda)\right).
\end{aligned}$$

Theorem 3.4. *Let $\Gamma_{[2]}^{(m,n)}$ be the subgroup of $\Gamma^{(m,n)}$ generated by the elements*

$$(S,(0,0)),\quad (T_*,(0,s))\quad \textit{and}\quad (I_2,(\lambda_0,\mu_0)),$$

where

$$T_*=\begin{pmatrix}1&2\\0&1\end{pmatrix}\quad \textit{and } s,\lambda_0,\mu_0\in\mathbb{Z}^{(m,n)}.$$

Let $\mathcal{M} = (\mathcal{M}_{kl})$ be a positive definite symmetric unimodular integral $m \times m$ matrix such that $\mathcal{M}\mathbb{Z}^{(m,n)} = \mathbb{Z}^{(m,n)}$. Then for $f, g \in \mathscr{S}\left(\mathbb{R}^{(m,n)}\right)$, the function

$$\Theta_f^{[\mathcal{M}]}(\tau, \phi; \lambda, \mu)\, \overline{\Theta_g^{[\mathcal{M}]}(\tau, \phi; \lambda, \mu)}$$

is invariant under the action of $\Gamma_{[2]}^{(m,n)}$ on $G^{(m,n)}$.

Proof. The proof follows directly from Theorem 3.1 (Jacobi 1), Theorem 3.2 (Jacobi 2) and Theorem 3.3 (Jacobi 3) because the left actions of the generators of $\Gamma_{[2]}^{(m,n)}$ are given by

$$((\tau, \phi), (\lambda, \mu)) \longmapsto \left(\left(-\frac{1}{\tau}, \phi + \arg\tau\right), (-\mu, \lambda)\right),$$
$$((\tau, \phi), (\lambda, \mu)) \longmapsto ((\tau + 2, \phi), (\lambda, s - 2\,\lambda + \mu))$$

and

$$((\tau, \phi), (\lambda, \mu)) \longmapsto ((\tau, \phi), (\lambda + \lambda_0, \mu + \mu_0)).$$

□

References

[1] R. Berndt and R. Schmidt, *Elements of theRepresentation Theory of the Jacobi Group*, Birkhäuser, 1998.

[2] M. Eichler and D. Zagier, *The Theory of Jacobi Forms*, Progress in Math., **55**, Birkhäuser, Boston, Basel and Stuttgart, 1985.

[3] E. Freitag, *Siegelsche Modulfunktionen*, Grundlehren de mathematischen Wissenschaften **55**, Springer-Verlag, Berlin-Heidelberg-New York (1983).

[4] M. Itoh, H. Ochiai and J.-H. Yang, *Invariant Differential Operators on the Siegel-Jacobi Space,* submitted (2015).

[5] G. Lion and M. Vergne, *The Weil representation, Maslov index and Theta seires*, Progress in Math., **6**, Birkhäuser, Boston, Basel and Stuttgart, 1980.

[6] J. Marklof, *Pair correlation densities of inhomogeneous quadratic forms*, Ann. of Math., **158** (2003), 419-471.

[7] D. Mumford, *Tata Lectures on Theta I,* Progress in Math. **28**, Boston-Basel-Stuttgart (1983).

[8] A. Pitale, *Jacobi Maass forms*, Abh. Math. Sem. Hamburg **79** (2009), 87–111.

[9] C. L. Siegel, *Indefinite quadratische Formen und Funnktionentheorie I and II*, Math. Ann. **124** (1951), 17–54 and Math. Ann. **124** (1952), 364–387 ; Gesammelte Abhandlungen, Band III, Springer-Verlag (1966), 105–142 and 154–177.

[10] A. Weil, *Sur certains groupes d'operateurs unitares*, Acta Math., **111** (1964), 143–211; Collected Papers (1964-1978), Vol. III, Springer-Verlag (1979), 1-69.

[11] J.-H. Yang, *Harmonic Analysis on the Quotient Spaces of Heisenberg Groups*, Nagoya Math. J., **123** (1991), 103–117.

[12] J.-H. Yang, *Harmonic Analysis on the Quotient Spaces of Heisenberg Groups II,* J. Number Theory, **49** (1) (1994), 63–72.

[13] J.-H. Yang, *A decomposition theorem on differential polynomials of theta functions of high level*, Japanese J. of Mathematics, the Mathematical Society of Japan, New Series, **22** (1) (1996), 37–49.

[14] J.-H. Yang, *Fock Representations of the Heisenberg Group $H_{\mathbb{R}}^{(g,h)}$*, J. Korean Math. Soc., **34**, no. 2 (1997), 345–370.

[15] J.-H. Yang, *Lattice Representations of the Heisenberg Group $H_{\mathbb{R}}^{(g,h)}$*, Math. Annalen, **317** (2000), 309–323.

[16] J.-H. Yang, *Heisenberg Group, Theta Functions and the Weil Representation,* Kyung Moon Sa, Seoul (2012).

[17] J.-H. Yang, *The Siegel-Jacobi Operator*, Abh. Math. Sem. Univ. Hamburg **63** (1993), 135–146.

[18] J.-H. Yang, *Remarks on Jacobi forms of higher degree*, Proc. of the 1993 Workshop on Automorphic Forms and Related Topics, the Pyungsan Institute for Mathematical Sciences, Seoul (1993), 33–58.
[19] J.-H. Yang, *Singular Jacobi forms,* Trans. of American Math. Soc. **347**, No. 6 (1995), 2041-2049.
[20] J.-H. Yang, *Construction of vector valued modular forms from Jacobi forms,* Canadian J. of Math. **47 (6)** (1995), 1329-1339.
[21] J.-H. Yang, *A note on a fundamental domain for Siegel-Jacobi space,* Houston Journal of Mathematics, Vol. **32**, No. 3 (2006), 701–712.
[22] J.-H. Yang, *Invariant metrics and Laplacians on Siegel-Jacobi space,* Journal of Number Theory, **127** (2007), 83–102.
[23] J.-H. Yang, *A partial Cayley transform of Siegel-Jacobi disk,* J. Korean Math. Soc. **45**, No. 3 (2008), 781-794.
[24] J.-H. Yang, *Invariant metrics and Laplacians on Siegel-Jacobi disk,* Chinese Annals of Mathematics, Vol. **31** B(1) (2010), 85-100.
[25] J.-H. Yang, *A Note on Maass-Jacobi Forms II,* Kyungpook Math. J. **53** (2013), 49-86.
[26] J.-H. Yang, Y.-H. Yong, S.-N. Huh, J.-H. Shin and G.-H. Min, *Sectional Curvatures of the Siegel-Jacobi Space,* Bull. Korean Math. Soc. **50** (2013), No. 3, pp. 787-799.
[27] J.-H. Yang, *Geometry and Arithmetic on the Siegel-Jacobi Space*, Geometry and Analysis on Manifolds, In Memory of Professor Shoshichi Kobayashi (edited by T. Ochiai, A. Weinstein et al), Progress in Mathematics, Volume **308**, Birkhäuser, Springer International Publishing AG Switzerland (2015), 275-325.
[28] J.-H. Yang, *Covariant maps for the Schrödinger-Weil representation,* Bull. Korean Math. Soc. **52** (2015), No. 2, pp. 627-647.
[29] C. Ziegler, *Jacobi Forms of Higher Degree*, Abh. Math. Sem. Hamburg **59** (1989), 191–224.

Department of Mathematics, Inha University, Incheon 22212, Korea
E-mail address: jhyang@inha.ac.kr